


The surveillance and control programme for infectious bovine rhinotracheitis (IBR) and infectious pustular vulvovaginitis (IPV) in Norway

Johan Åkerstedt

Attila Tarpai

Tormod Mørk



*Scientific editors Hege Hellberg and Ståle Sviland
Norwegian Veterinary Institute*



**Surveillance and control programmes for terrestrial and aquatic animals in Norway.
Annual report 2010**

Editors

Ståle Sviland and Hege Hellberg

Publisher

Norwegian Veterinary Institute
PO Box 750 Sentrum
N-0106 Oslo
Norway

Fax: + 47 23 21 60 01

Tel: + 47 23 21 60 00

E-mail: postmottak@vetinst.no

www.vetinst.no

Front page photo: Hanne Mari Jordsmyr

ISSN 1503-1454

Example of citation:

Åkerstedt J, Tarpai A, Mørk T. The surveillance and control programme for infectious bovine rhinotracheitis (IBR) and infectious pustular vulvovaginitis (IPV) in Norway. In: Sviland S, Hellberg H (editors). Surveillance and control programmes for terrestrial and aquatic animals in Norway. Annual report 2010. Oslo: Norwegian Veterinary Institute; 2011. ISSN 1503-1454.

© Norwegian Veterinary Institute

Any use of the present data should include specific reference to this report.

The surveillance and control programme for infectious bovine rhinotracheitis (IBR) and infectious pustular vulvovaginitis (IPV) in Norway

Johan Åkerstedt, Attila Tarpai, Tormod Mørk

All milk and blood samples tested in 2010 were negative for antibodies against bovine herpes virus (BHV-1).

Introduction

Infectious bovine rhinotracheitis and infectious pustular vulvovaginitis (IBR/IPV) is a disease caused by bovine herpesvirus 1 (BoHV-1). The virus affects the upper respiratory tract of cattle causing purulent nasal discharge, hyperaemia of the muzzle and conjunctivitis. Signs of general illness are fever, depression, reduced appetite, abortions and reduced milk yield. The virus may also infect the genital tract and cause pustular vulvovaginitis and balanoposthitis. IBR/IPV is classified as list B disease in Norway and is notifiable to the Office International des Epizooties.

Norway has not experienced clinical outbreaks of infectious pustular vulvovaginitis since the early 1960s, when two outbreaks were diagnosed in cattle. In 1993, animals in one single herd were found seropositive after primary testing of bulk milk samples. Clinical signs of IBR/IPV were not recorded and all animals on the farm were slaughtered. Attempts to isolate the virus from organ samples gave negative results. Contact herds and dairy herds in the same region were found serologically negative (1). Likewise, red deer that were shot in the neighbourhood during the hunting season the same year were found seronegative. After this incident, IBR/IPV virus infection has not been demonstrated in Norway.

EFTA Surveillance Authority (ESA) has recognised Norway as free from IBR since 1994. Decisions concerning the additional guarantees relating to IBR/IPV for bovines destined for Norway are described in ESA Decision 74/94/COL. Maintenance of the ESA Decisions accepting the IBR-free status of Norway requires annual reports of the surveillance of the disease.

The Norwegian Food Safety Authority was responsible for carrying out the surveillance and control programme for IBR/IPV. The Norwegian Veterinary Institute was in charge of planning the programme, collecting the bulk milk samples from the dairies and performing the tests. Blood samples from beef herds were collected by inspectors from the Norwegian Food Safety Authority.

Aim

The aim of the surveillance and control programme for IBR/IPV was to document freedom from the infection in Norway according to the demands in ESA Decision 74/94/COL with amendments, and to contribute to the maintenance of this favourable situation.

Material and methods

The surveillance of cattle for IBR/IPV in 2010 included both dairy and beef herds. Bulk milk samples from the dairy herds were provided by the dairies. From the beef herds, individual blood samples were collected on the farms from maximum ten animals older than 24 months.

The target population of dairy herds consisted of all cattle herds delivering milk to dairies during the sampling period. In 2010, bulk milk samples from 1,265 randomly sampled dairy herds were tested. The group of beef herds to be sampled was based on a register of all beef herds receiving governmental support according to recordings of July 2010. A total of 4,020 individual blood samples from 507 beef herds were analysed in herd pools. The sampled herds represented 11.6% of the Norwegian cattle herds (Table 1).

The number of herds in the surveillance and control programme for IBR/IPV in 2010 is given in Table 1. All samples were tested for antibodies against bovine herpes virus 1 (BHV-1) using a commercial indirect enzyme-linked immunosorbent assay (ELISA; Svanova Biotech AB, Uppsala, Sweden) at the Norwegian Veterinary Institute in Sandnes. In case of any positive or dubious results, a serum neutralization test would be preformed.

Table 1. Numbers of dairy herds and beef herds within the frame of the Norwegian surveillance and control programme for IBR/IPV in 2010

Herd category	Total no. of cattle herds*	No. of herds tested	% tested of the total no. of herds
Dairy herds**	11,150	1,265	11.3
Beef herds	4,200	507	12.1
Total	15,350	1,772	11.6

* Based on data from the Register of production subsidies as of 31 July 2010.

** Cattle herds delivering milk to dairies.

Results

All bulk milk samples and blood samples tested in 2010 were negative for antibodies against BHV-1. Table 2 shows the results of the testing during the period from 1993 to 2010.

Table 2. Samples in the surveillance and control programme for IBR/IPV in the Norwegian bovine population during the period 1993-2010

Year	Dairy herds	Beef herds		No. of positive samples
	No. of bulk milk samples tested	No. of beef herds sampled	No. of individuals tested	
1993	26,642	0	0	1
1994	24,832	1,430	5,954	0
1995	25,131	1,532	9,354	0
1996	2,863	303	1,523	0
1997	2,654	2,214	16,741	0
1998	2,816	2,191	17,095	0
1999	2,930	2,382	18,274	0
2000	1,590	340	2,892	0
2001	2,564	434	3,453	0
2002	2,308	462	3,693	0
2003	1,845	449	3,901	0
2004	1,573	402	3,364	0
2005	1,919	484	4,766	0
2006	1,673	479	4,624	0
2007	1,575	412	4,241	0
2008	1,422	444	4,616	0
2009	1,315	435	5,048	0
2010	1,265	507	4,020	0

Discussion

Since 2008, a commercial indirect ELISA has replaced the previously used blocking ELISA (2). The latter method had been evaluated in a retrospective analysis using a simulation model (3). However, after participation in a proficiency testing scheme organized by the Veterinary Laboratories Agency Weybridge, New Haw, England, the indirect ELISA was found better suited for testing bulk milk specifically, and this method was preferred.

In addition to the surveillance programme, all breeding bull candidates are serologically tested before entering the breeding centres, and all breeding bulls are subject to a compulsory test each year.

The results of the programme since 1993 strongly indicate that the Norwegian cattle population is free from IBR/IPV-infection (4).

References

1. Tharaldsen J, Krogsrud J, Ødegaard Ø. Påvist besetningsinfeksjon med bovint herpesvirus 1 (BHV-1) [Herd infection with bovine herpes virus (BHV-1) detected, No]. *Nor Vet Tidsskr* 1993; 105: 363-4.
2. Nylin B, Strøger U, Rønsholt L. A retrospective evaluation of a bovine herpes virus-1 (BHV-1) antibody ELISA on bulk-tank milk samples for classification of the BHV-1 status of Danish dairy herds. *Prev Vet Med* 2000; 47: 91-105.
3. Paisley LG, Tharaldsen J, Jarp J. A retrospective analysis of the infectious rhinotracheitis (bovine herpes virus-1) surveillance program using Monte Carlo simulation models. *Prev Vet Med* 2001; 50: 109-25.
4. Åkerstedt J, Tarpai A, Mørk T. The surveillance and control programme for infectious bovine rhinotracheitis (IBR) and infectious pustular vulvovaginitis (IPV) in Norway. Annual report 2009. In: Karlsson AC, Jordsmyr HM, Hellberg H, Sviland s (editors). *Surveillance and control programmes for terrestrial and aquatic animals in Norway*. Oslo: National Veterinary Institute; 2010.

The Norwegian Veterinary Institute (NVI) is a nationwide research institute in the fields of animal health, fish health, and food safety. The primary mission of the NVI is to give research-based independent advisory support to ministries and governing authorities. Preparedness, diagnostics, surveillance, reference functions, risk assessments, and advisory and educational functions are the most important areas of operation.

The Norwegian Veterinary Institute has its main laboratory in Oslo, with regional laboratories in Sandnes, Bergen, Trondheim, Harstad og Tromsø, with about 360 employees in total.

www.vetinst.no



Veterinærinstituttet
Norwegian Veterinary Institute

The Norwegian Food Safety Authority (NFSA) is a governmental body whose aim is to ensure through regulations and controls that food and drinking water are as safe and healthy as possible for consumers and to promote plant, fish and animal health and ethical farming of fish and animals. We encourage environmentally friendly production and we also regulate and control cosmetics, veterinary medicines and animal health personnel. The NFSA drafts and provides information on legislation, performs risk-based inspections, monitors food safety, plant, fish and animal health, draws up contingency plans and provides updates on developments in our field of competence.

The NFSA comprises three administrative levels, and has some 1300 employees.

The NFSA advises and reports to the Ministry of Agriculture and Food, the Ministry of Fisheries and Coastal Affairs and the Ministry of Health and Care Services.

www.mattilsynet.no

